

Preprint, arXiv:0911.4433v7

ARITHMETIC THEORY OF HARMONIC NUMBERS (II)

ZHI-WEI SUN¹ AND LI-LU ZHAO²

¹Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
zwsun@nju.edu.cn
<http://math.nju.edu.cn/~zwsun>

²Department of Mathematics
The University of Hong Kong
Pokfulam Road, Hong Kong
zhaolilu@gmail.com

ABSTRACT. For $k = 0, 1, 2, \dots$ let H_k denote the harmonic number $\sum_{0 < j \leq k} 1/j$. In this paper we establish some new congruences involving harmonic numbers. For example, we show that for any prime $p > 3$ we have

$$\sum_{k=1}^{p-1} \frac{H_k}{k2^k} \equiv \frac{7}{24} p B_{p-3} \pmod{p^2}, \quad \sum_{k=1}^{p-1} \frac{H_{k,2}}{k2^k} \equiv -\frac{3}{8} B_{p-3} \pmod{p},$$

and

$$\sum_{k=1}^{p-1} \frac{H_{k,2n}^2}{k^{2n}} \equiv \frac{\binom{6n+1}{2n-1} + n}{6n+1} p B_{p-1-6n} \pmod{p^2}$$

for any positive integer $n < (p-1)/6$, where $B_0, B_1, B_2, \dots$ are Bernoulli numbers, and $H_{k,m} := \sum_{0 < j \leq k} 1/j^m$.

1. INTRODUCTION

Recall that harmonic numbers are those

$$H_n := \sum_{0 < k \leq n} \frac{1}{k} \quad (n \in \mathbb{N} = \{0, 1, 2, \dots\}).$$

2010 *Mathematics Subject Classification*. Primary 11A07, 11B68; Secondary 05A19, 11B75.

Keywords. Harmonic numbers, congruences, Bernoulli numbers.

The initial version of this paper was posted to arXiv on August 5, 2011.

The first author is supported by the National Natural Science Foundation (grant 11171140) of China.

They play important roles in mathematics. In 1862 J. Wolstenholme [W] showed that $H_{p-1} \equiv 0 \pmod{p^2}$ for any prime $p > 3$.

In [Su] the first author investigated arithmetic properties of harmonic numbers systematically. For example, he proved that for any prime $p > 5$ we have

$$\sum_{k=1}^{p-1} \frac{H_k}{k2^k} \equiv \sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv 0 \pmod{p}.$$

For $m \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$, harmonic numbers of order m are defined by

$$H_{n,m} := \sum_{0 < k \leq n} \frac{1}{k^m} \quad (n \in \mathbb{N}).$$

It is known that

$$\sum_{k=1}^{\infty} \frac{H_k}{k2^k} = \frac{\pi^2}{12} \quad (\text{S. W. Coffman [C], 1987})$$

and

$$\sum_{k=1}^{\infty} \frac{H_{k,2}}{k2^k} = \frac{5}{8}\zeta(3) \quad (\text{B. Cloitre, 2004}).$$

Both identities can be found in [SW].

Our first theorem is as follows.

Theorem 1.1. *For any prime $p > 3$, we have*

$$\sum_{k=0}^{p-1} \frac{H_k}{k2^k} \equiv \frac{7}{24}pB_{p-3} \pmod{p^2} \quad (1.1)$$

and

$$\sum_{k=1}^{p-1} \frac{H_{k,2}}{k2^k} \equiv -\frac{3}{8}B_{p-3} \pmod{p}, \quad (1.2)$$

where $B_0, B_1, B_2, \dots$ are Bernoulli numbers.

Remark 1.1. (1.1) confirms the first part of [Su, Conjecture 1.1]. The second part of [Su, Conjecture 1.1] states that $\sum_{k=1}^{p-1} H_k^2/k^2 \equiv \frac{4}{5}pB_{p-5} \pmod{p^2}$ for any prime $p > 3$; this was confirmed by R. Meštrović [M] quite recently.

Our second theorem confirms the second conjecture of [Su].

Theorem 1.2 ([Su, Conjecture 1.2]). *Let p be an odd prime and let n be a positive integer with $p-1 \nmid 6n$. Then*

$$\sum_{k=1}^{p-1} \frac{H_{k,2n}^2}{k^{2n}} \equiv 0 \pmod{p}. \quad (1.3)$$

Furthermore, when $p > 6n + 1$ we have

$$\sum_{k=1}^{p-1} \frac{H_{k,2n}^2}{k^{2n}} \equiv \frac{s(n)}{6n+1} p B_{p-1-6n} \pmod{p^2}, \quad (1.4)$$

where

$$s(n) = \binom{6n+1}{2n-1} + n.$$

Remark 1.2. We give here four initial values of the integer sequence $\{s(n)\}_{n \geq 1}$:

$$s(1) = 8, \quad s(2) = 288, \quad s(3) = 11631, \quad s(4) = 480704.$$

We will show Theorems 1.1 and 1.2 in Sections 2 and 3 respectively.

2. PROOF OF THEOREM 1.1

Lemma 2.1. *Let $p > 3$ be a prime. Then*

$$\sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} \equiv \frac{p}{2} B_{p-3} \pmod{p^2}, \quad \sum_{k=1}^{p-1} \frac{(-1)^k}{k^3} \equiv -\frac{B_{p-3}}{2} \pmod{p}, \quad (2.1)$$

and

$$\sum_{k=1}^{p-1} \frac{H_k}{k} \equiv \frac{p}{3} B_{p-3} \pmod{p^2} \quad \text{and} \quad \sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} H_k \equiv -\frac{B_{p-3}}{4} \pmod{p}. \quad (2.2)$$

Proof. It is known that (cf. [S, Corollaries 5.1 and 5.2])

$$\sum_{k=1}^{p-1} \frac{1}{k^2} \equiv \frac{2}{3} p B_{p-3} \pmod{p^2}, \quad \sum_{k=1}^{p-1} \frac{1}{k^3} \equiv \frac{3}{4} p B_{p-4} \equiv -p \delta_{p,5} \pmod{p^2},$$

and

$$\sum_{k=1}^{(p-1)/2} \frac{1}{k^2} \equiv \frac{7}{3} p B_{p-3} \pmod{p^2} \quad \text{and} \quad \sum_{k=1}^{(p-1)/2} \frac{1}{k^3} \equiv -2 B_{p-3} \pmod{p}.$$

Thus

$$\begin{aligned}\sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} &= \sum_{k=1}^{p-1} \frac{1+(-1)^k}{k^2} - \sum_{k=1}^{p-1} \frac{1}{k^2} = \frac{1}{2} H_{(p-1)/2,2} - H_{p-1,2} \\ &\equiv \frac{7}{6} p B_{p-3} - \frac{2}{3} p B_{p-3} \equiv \frac{p}{2} B_{p-3} \pmod{p^2}\end{aligned}$$

and

$$\begin{aligned}\sum_{k=1}^{p-1} \frac{(-1)^k}{k^3} &= \sum_{k=1}^{p-1} \frac{1+(-1)^k}{k^3} - \sum_{k=1}^{p-1} \frac{1}{k^3} \\ &= \frac{1}{4} H_{(p-1)/2,3} - H_{p-1,3} \equiv \frac{-2B_{p-3}}{4} \pmod{p}.\end{aligned}$$

Therefore (2.1) holds.

By the proof of [S, Theorem 6.1],

$$\sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk} \equiv -\frac{p}{3} B_{p-3} \pmod{p^2}.$$

So we have

$$\sum_{k=1}^{p-1} \frac{H_k}{k} = \sum_{k=1}^{p-1} \frac{1}{k^2} + \sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk} \equiv \frac{2}{3} p B_{p-3} - \frac{p}{3} B_{p-3} = \frac{p}{3} B_{p-3} \pmod{p^2}.$$

This proves the first congruence in (2.2).

Now we prove the second congruence in (2.2). Since

$$H_{p-k} = H_{p-1} - \sum_{j=1}^{k-1} \frac{1}{p-j} \equiv H_{k-1} = H_k - \frac{1}{k} \pmod{p}$$

for all $k = 1, \dots, p-1$, we have

$$\sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} H_k = \sum_{k=1}^{p-1} \frac{(-1)^{p-k}}{(p-k)^2} H_{p-k} \equiv - \sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} \left(H_k - \frac{1}{k} \right) \pmod{p}$$

and hence

$$\sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} H_k \equiv \frac{1}{2} \sum_{k=1}^{p-1} \frac{(-1)^k}{k^3} \equiv -\frac{B_{p-3}}{4} \pmod{p}.$$

The proof of Lemma 2.1 is now complete. $\square$

Lemma 2.2. (i) *For any positive integers k and m we have*

$$\sum_{n=1}^m \binom{n-1}{k-1} = \binom{m}{k}. \quad (2.3)$$

(ii) *For each $n = 1, 2, 3, \dots$ we have*

$$\sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k-1}}{k} H_k = H_{n,2}. \quad (2.4)$$

Proof. (2.3) is well known (cf. [G, (1.5)]) and it can be easily proved by induction on m .

(2.4) is also known, but it is difficult to give a reference. Here we prove it by induction. Clearly (2.4) holds for $n = 1$. Assume that (2.4) holds for a fixed positive integer n . Then

$$\begin{aligned} \sum_{k=1}^{n+1} \binom{n+1}{k} \frac{(-1)^{k-1}}{k} H_k &= \sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k-1}}{k} H_k + \sum_{k=1}^{n+1} \binom{n}{k-1} \frac{(-1)^{k-1}}{k} H_k \\ &= H_{n,2} + \frac{1}{n+1} \sum_{k=0}^{n+1} \binom{n+1}{k} (-1)^{k-1} H_k. \end{aligned}$$

Note that

$$\begin{aligned} &\sum_{k=0}^{n+1} \binom{n+1}{k} (-1)^{k-1} H_k \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^{k-1} H_k + \sum_{k=1}^{n+1} \binom{n}{k-1} (-1)^{k-1} \left(H_{k-1} + \frac{1}{k} \right) \\ &= \sum_{k=1}^{n+1} \binom{n}{k-1} \frac{(-1)^{k-1}}{k} = -\frac{1}{n+1} \sum_{k=1}^{n+1} \binom{n+1}{k} (-1)^k = \frac{1}{n+1}. \end{aligned}$$

So

$$\sum_{k=1}^{n+1} \binom{n+1}{k} \frac{(-1)^{k-1}}{k} H_k = H_{n,2} + \frac{1}{n+1} \cdot \frac{1}{n+1} = H_{n+1,2}$$

as desired. This concludes the proof. $\square$

Lemma 2.3. *Let $p > 3$ be a prime. Then*

$$\sum_{1 \leq j \leq k \leq p-1} \frac{2^j(j+k)}{j^2 k^2} \equiv \sum_{k=1}^{p-1} \frac{(-1)^k}{k^3} \pmod{p}. \quad (2.5)$$

Proof. Observe that

$$\begin{aligned}
& \sum_{1 \leq i \leq j \leq k \leq p-1} \frac{2^i}{ijk} - \sum_{1 \leq i < j < k \leq p-1} \frac{2^i}{ijk} \\
&= \sum_{1 \leq j \leq k \leq p-1} \frac{2^j}{j^2 k} + \sum_{1 \leq i \leq j \leq p-1} \frac{2^i}{ij^2} - \sum_{k=1}^{p-1} \frac{2^k}{k^3} \\
&= \sum_{1 \leq j \leq k \leq p-1} \left(\frac{2^j}{j^2 k} + \frac{2^j}{jk^2} \right) - \sum_{k=1}^{p-1} \frac{2^k}{k^3}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
& 2 \sum_{1 \leq i \leq j \leq k \leq p-1} \frac{(-1)^i}{ijk} - 2 \sum_{1 \leq i < j < k \leq p-1} \frac{(-1)^i}{ijk} - 2 \sum_{k=1}^{p-1} \frac{(-1)^k}{k^3} \\
&= 2 \sum_{1 \leq j < k \leq p-1} \left(\frac{(-1)^j}{j^2 k} + \frac{(-1)^j}{jk^2} \right) \\
&\equiv \sum_{1 \leq j < k \leq p-1} \left(\frac{(-1)^j}{j^2 k} + \frac{(-1)^j}{jk^2} + \frac{(-1)^{p-j}}{(p-j)^2(p-k)} + \frac{(-1)^{p-j}}{(p-j)(p-k)^2} \right) \\
&= \sum_{1 \leq j < k \leq p-1} \frac{(-1)^j}{j^2 k} + \sum_{1 \leq k < j \leq p-1} \frac{(-1)^j}{j^2 k} \\
&\quad + \sum_{1 \leq j < k \leq p-1} \frac{(-1)^j}{jk^2} + \sum_{1 \leq k < j \leq p-1} \frac{(-1)^j}{jk^2} \\
&= H_{p-1} \sum_{j=1}^{p-1} \frac{(-1)^j}{j^2} + H_{p-1,2} \sum_{j=1}^{p-1} \frac{(-1)^j}{j} - 2 \sum_{j=1}^{p-1} \frac{(-1)^j}{j^3} \pmod{p}.
\end{aligned}$$

Thus, with the help of $H_{p-1} \equiv H_{p-1,2} \equiv 0 \pmod{p}$, we have

$$\sum_{1 \leq i \leq j \leq k \leq p-1} \frac{(-1)^i}{ijk} \equiv \sum_{1 \leq i < j < k \leq p-1} \frac{(-1)^i}{ijk} \pmod{p}.$$

By [ZS, Theorem 1.2],

$$\sum_{1 \leq i < j < k \leq p-1} \frac{(1-x)^i}{ijk} \equiv \sum_{1 \leq i < j < k \leq p-1} \frac{x^i}{ijk} \pmod{p}.$$

So, in view of the above, we have

$$\begin{aligned}
\sum_{1 \leq i \leq j \leq k \leq p-1} \frac{(-1)^i}{ijk} &\equiv \sum_{1 \leq i < j < k \leq p-1} \frac{2^i}{ijk} \\
&\equiv \sum_{1 \leq i \leq j \leq k \leq p-1} \frac{2^i}{ijk} + \sum_{k=1}^{p-1} \frac{2^k}{k^3} - \sum_{1 \leq j < k \leq p-1} \frac{2^j(j+k)}{j^2 k^2} \pmod{p}.
\end{aligned}$$

Thus, it remains to show that

$$\sum_{1 \leq i \leq j \leq k \leq p-1} \frac{2^i - (-1)^i}{ijk} \equiv \sum_{k=1}^{p-1} \frac{(-1)^k - 2^k}{k^3} \pmod{p}. \quad (2.6)$$

With the help of Lemma 2.2, we have

$$\begin{aligned} \sum_{1 \leq i \leq j \leq k \leq p-1} \frac{2^i - (-1)^i}{ijk} &= \sum_{1 \leq i \leq j \leq k \leq p-1} \frac{1}{ijk} \sum_{r=0}^i (1 - (-2)^r) \binom{i}{r} \\ &= \sum_{r=1}^{p-1} \frac{1 - (-2)^r}{r} \sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk} \sum_{i=1}^j \binom{i-1}{r-1} \\ &= \sum_{r=1}^{p-1} \frac{1 - (-2)^r}{r} \sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk} \binom{j}{r} \\ &= \sum_{r=1}^{p-1} \frac{1 - (-2)^r}{r^2} \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=1}^k \binom{j-1}{r-1} \\ &= \sum_{r=1}^{p-1} \frac{1 - (-2)^r}{r^2} \sum_{k=1}^{p-1} \frac{1}{k} \binom{k}{r} = \sum_{r=1}^{p-1} \frac{1 - (-2)^r}{r^3} \sum_{k=1}^{p-1} \binom{k-1}{r-1} \\ &= \sum_{r=1}^{p-1} \frac{1 - (-2)^r}{r^3} \binom{p-1}{r} \equiv \sum_{r=1}^{p-1} \frac{(-1)^r - 2^r}{r^3} \pmod{p}. \end{aligned}$$

This proves the desired (2.6). $\square$

Proof of Theorem 1.1. (i) We prove (1.2) first. In view of (2.4), we have

$$\begin{aligned} \sum_{n=1}^{p-1} \frac{H_{n,2}}{n2^n} &= \sum_{n=1}^{p-1} \frac{1}{n2^n} \sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k-1}}{k} H_k \\ &= \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} H_k \sum_{n=k}^{p-1} \frac{1}{n2^n} \binom{n}{k} = \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k^2 2^k} H_k \sum_{n=k}^{p-1} \binom{n-1}{k-1} \frac{1}{2^{n-k}} \\ &= \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k^2 2^k} H_k \sum_{j=0}^{p-1-k} \binom{k+j-1}{j} \frac{1}{2^j} \\ &= \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k^2 2^k} H_k \sum_{j=0}^{p-1-k} \binom{-k}{j} \frac{1}{(-2)^j} \end{aligned}$$

and hence

$$\begin{aligned}
\sum_{n=1}^{p-1} \frac{H_{n,2}}{n2^n} &\equiv \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k^2 2^k} H_k \sum_{j=0}^{p-1-k} \binom{p-k}{j} \frac{1}{(-2)^j} \\
&= \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k^2 2^k} H_k \frac{1 + (-1)^k}{2^{p-k}} \\
&\equiv -\frac{1}{2} \sum_{k=1}^{p-1} \frac{H_k}{k^2} (1 + (-1)^k) \pmod{p}.
\end{aligned}$$

Note that

$$\sum_{k=1}^{p-1} \frac{H_k}{k^2} \equiv B_{p-3} \pmod{p} \quad \text{and} \quad \sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} H_k \equiv -\frac{B_{p-3}}{4} \pmod{p}$$

by [ST, (5.4)] and (2.2) respectively. So we get

$$\sum_{n=1}^{p-1} \frac{H_{n,2}}{n2^n} \equiv -\frac{1}{2} \left(B_{p-3} - \frac{B_{p-3}}{4} \right) = -\frac{3}{8} B_{p-3} \pmod{p}.$$

(ii) Now we show (1.1). Observe that

$$\begin{aligned}
\sum_{k=1}^{p-1} \frac{H_k}{k2^k} &= \sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk2^k} = \sum_{1 \leq j \leq k \leq p-1} \frac{1}{(p-k)(p-j)2^{p-j}} \\
&= \sum_{1 \leq j \leq k \leq p-1} \frac{2^{j-p}(p+j)(p+k)}{(p^2-j^2)(p^2-k^2)} \\
&\equiv \sum_{1 \leq j \leq k \leq p-1} \frac{2^{j-p}(jk + p(j+k))}{j^2 k^2} \\
&= 2^{-p} \sum_{1 \leq j \leq k \leq p-1} \frac{2^j}{jk} + \frac{p}{2} \sum_{1 \leq j \leq k \leq p-1} \frac{2^j(j+k)}{j^2 k^2} \pmod{p^2}.
\end{aligned}$$

In view of Lemmas 2.2 and 2.1,

$$\begin{aligned}
\sum_{1 \leq j \leq k \leq p-1} \frac{2^j - 1}{jk} &= \sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk} \sum_{i=1}^j \binom{j}{i} = \sum_{i=1}^{p-1} \frac{1}{i} \sum_{k=1}^{p-1} \frac{1}{k} \sum_{j=1}^k \binom{j-1}{i-1} \\
&= \sum_{i=1}^{p-1} \frac{1}{i} \sum_{k=1}^{p-1} \frac{1}{k} \binom{k}{i} = \sum_{i=1}^{p-1} \frac{1}{i^2} \sum_{k=1}^{p-1} \binom{k-1}{i-1} \\
&= \sum_{i=1}^{p-1} \frac{1}{i^2} \binom{p-1}{i} = \sum_{i=1}^{p-1} \frac{(-1)^i}{i^2} \prod_{r=1}^i \left(1 - \frac{p}{r} \right) \\
&\equiv \sum_{i=1}^{p-1} \frac{(-1)^i (1 - pH_i)}{i^2} \equiv \frac{p}{2} B_{p-3} - p \left(-\frac{B_{p-3}}{4} \right) \pmod{p^2}.
\end{aligned}$$

Note that

$$\sum_{1 \leq j \leq k \leq p-1} \frac{1}{jk} = \sum_{k=1}^{p-1} \frac{H_k}{k} \equiv \frac{p}{3} B_{p-3} \pmod{p^2}$$

by (2.2). Combining the above with (2.5), we finally obtain that

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{H_k}{k2^k} &\equiv 2^{-p} \left(\frac{3}{4} p B_{p-3} + \frac{p}{3} B_{p-3} \right) + \frac{p}{2} \sum_{k=1}^{p-1} \frac{(-1)^k}{k^3} \\ &\equiv \frac{13}{24} p B_{p-3} + \frac{p}{2} \left(-\frac{B_{p-3}}{2} \right) = \frac{7}{24} p B_{p-3} \pmod{p^2} \quad (\text{by (2.1)}). \end{aligned}$$

This concludes the proof. $\square$

3. PROOF OF THEOREM 1.2

Lemma 3.1. *Let $p > 3$ be a prime and let m be a positive even integer with $p-1 \nmid 3m$. Then*

$$\sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^m k^{2m}} + \frac{1}{j^{2m} k^m} \right) \equiv 0 \pmod{p}. \quad (3.1)$$

Moreover, if $p > 3m+1$, then

$$\sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^m k^{2m}} + \frac{1}{j^{2m} k^m} \right) \equiv -p \frac{3m}{3m+1} B_{p-1-3m} \pmod{p^2}. \quad (3.2)$$

Proof. It is well-known that

$$\sum_{k=1}^{p-1} \frac{1}{k^n} \equiv 0 \pmod{p} \quad \text{for any integer } n \not\equiv 0 \pmod{p-1}.$$

Also,

$$\sum_{k=1}^{p-1} \frac{1}{k^n} \equiv \frac{pn}{n+1} B_{p-1-n} \pmod{p^2} \quad \text{for } n = 2, \dots, p-3$$

(see, e.g., [S, Corollary 5.1]). Thus

$$\sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^m k^{2m}} + \frac{1}{j^{2m} k^m} \right) = \sum_{j=1}^{p-1} \frac{1}{j^m} \sum_{k=1}^{p-1} \frac{1}{k^{2m}} - \sum_{k=1}^{p-1} \frac{1}{k^{3m}} \equiv 0 \pmod{p}.$$

Moreover, we have (3.2) if $p > 3m+1$. $\square$

Lemma 3.2. *Let $p > 3$ be a prime and let m be a positive even integer. Then*

$$\sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^m k^{2m}} - \frac{1}{j^{2m} k^m} \right) \equiv 0 \pmod{p}. \quad (3.3)$$

Moreover, if $p > 3m + 1$ then

$$\sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^m k^{2m}} - \frac{1}{j^{2m} k^m} \right) \equiv \frac{pm \binom{3m}{m} B_{p-1-3m}}{(m+1)(2m+1)} \pmod{p^2}. \quad (3.4)$$

Proof. As m is even, we have

$$\begin{aligned} \sum_{1 \leq j < k \leq p-1} \frac{1}{j^m k^{2m}} &= \sum_{1 \leq j < k \leq p-1} \frac{1}{(p-k)^m (p-j)^{2m}} \\ &\equiv \sum_{1 \leq j < k \leq p-1} \frac{1}{j^{2m} k^m} \pmod{p}. \end{aligned}$$

Now suppose that $p > 3m + 1$. Then

$$\begin{aligned} \sum_{1 \leq j < k \leq p-1} \frac{1}{j^m k^{2m}} &= \sum_{1 \leq j < k \leq p-1} \frac{(p+k)^m (p+j)^{2m}}{(p^2-k^2)^m (p^2-j^2)^{2m}} \\ &\equiv \sum_{1 \leq j < k \leq p-1} \frac{(k^m + pmk^{m-1})(j^{2m} + p2mj^{2m-1})}{j^{4m} k^{2m}} \\ &\equiv \sum_{1 \leq i < j \leq p-1} \frac{1}{j^{2m} k^m} + pm \sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^{2m} k^{m+1}} + \frac{2}{j^{2m+1} k^m} \right) \pmod{p^2}. \end{aligned}$$

So, (3.4) is reduced to

$$\sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^{2m} k^{m+1}} + \frac{2}{j^{2m+1} k^m} \right) \equiv \frac{\binom{3m}{m} B_{p-1-3m}}{(m+1)(2m+1)} \pmod{p}. \quad (3.5)$$

Recall that for any integer n we have

$$\sum_{k=1}^{p-1} k^n \equiv \begin{cases} p-1 \pmod{p} & \text{if } p-1 \mid n, \\ 0 \pmod{p} & \text{if } p-1 \nmid n. \end{cases}$$

(See, e.g., [IR, p.235].) Also,

$$\sum_{j=0}^{k-1} j^n = \frac{1}{n+1} \sum_{j=0}^n \binom{n+1}{j} B_j k^{n+1-j}$$

for any $k = 1, 2, 3, \dots$ and $n = 0, 1, 2, \dots$ (See, e.g., [IR].) Therefore

$$\begin{aligned}
& \sum_{1 \leq j < k \leq p-1} \frac{1}{j^{2m} k^{m+1}} \\
& \equiv \sum_{k=1}^{p-1} \frac{1}{k^{m+1}} \sum_{j=0}^{k-1} j^{p-1-2m} = \sum_{k=1}^{p-1} \frac{1}{k^{m+1}(p-2m)} \sum_{j=0}^{p-1-2m} \binom{p-2m}{j} B_j k^{p-2m-j} \\
& \equiv -\frac{1}{2m} \sum_{j=0}^{p-1-2m} \binom{p-2m}{j} B_j \sum_{k=1}^{p-1} k^{p-1-3m-j} \\
& \equiv \frac{1}{2m} \sum_{\substack{j=0 \\ p-1|j+3m}}^{p-1-2m} \binom{p-2m}{j} B_j = \frac{1}{2m} \binom{p-2m}{m+1} B_{p-1-3m} \\
& \equiv \frac{1}{2m} \binom{-2m}{m+1} B_{p-1-3m} = \frac{(-1)^{m+1}}{2m} \binom{3m}{m+1} B_{p-1-3m} \pmod{p}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \sum_{1 \leq j < k \leq p-1} \frac{1}{j^{2m+1} k^m} \\
& \equiv \sum_{k=1}^{p-1} \frac{1}{k^m} \sum_{j=0}^{k-1} j^{p-2-2m} = \sum_{k=1}^{p-1} \frac{1}{k^m(p-1-2m)} \sum_{j=0}^{p-2-2m} \binom{p-1-2m}{j} B_j k^{p-1-2m-j} \\
& \equiv -\frac{1}{2m+1} \sum_{j=0}^{p-2-2m} \binom{p-1-2m}{j} B_j \sum_{k=1}^{p-1} k^{p-1-3m-j} \\
& \equiv \frac{1}{2m+1} \sum_{\substack{j=0 \\ p-1|j+3m}}^{p-2-2m} \binom{p-1-2m}{j} B_j = \frac{1}{2m+1} \binom{p-1-2m}{m} B_{p-1-3m} \\
& \equiv \frac{1}{2m+1} \binom{-1-2m}{m+1} B_{p-1-3m} = \frac{(-1)^m}{2m+1} \binom{3m}{m} B_{p-1-3m} \pmod{p}.
\end{aligned}$$

Therefore

$$\begin{aligned}
& \sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^{2m} k^{m+1}} + \frac{2}{j^{2m+1} k^m} \right) \\
& \equiv \left(\frac{(-1)^{m+1}}{2m} \binom{3m}{m+1} + 2 \frac{(-1)^m}{2m+1} \binom{3m}{m} \right) B_{p-1-3m} \\
& = \frac{(-1)^m}{(m+1)(2m+1)} \binom{3m}{m} B_{p-1-3m} \pmod{p}.
\end{aligned}$$

So (3.5) holds as m is even. We are done. $\square$

Proof of Theorem 1.2. Let $m = 2n$. Clearly

$$\begin{aligned}
\sum_{k=1}^{p-1} \frac{H_{k,m}^2}{k^m} &= \sum_{k=1}^{p-1} \frac{1}{k^m} \left(\sum_{j=1}^k \frac{1}{j^m} \right)^2 \\
&= \sum_{k=1}^{p-1} \frac{1}{k^m} \left(\sum_{j=1}^k \frac{1}{j^m} + 2 \sum_{1 \leq i < j \leq k} \frac{1}{i^m j^{2m}} \right) \\
&= H_{p-1,3m} + \sum_{1 \leq j < k \leq p-1} \frac{1}{j^{2m} k^m} + 2 \sum_{1 \leq i < j \leq p-1} \frac{1}{i^m j^{2m}} \\
&\quad + 2 \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^m j^m k^m}
\end{aligned}$$

and

$$\begin{aligned}
H_{p-1,m}^3 &= \sum_{i=1}^{p-1} \frac{1}{i^m} \left(\sum_{k=1}^{p-1} \frac{1}{k^{2m}} + 2 \sum_{1 \leq j < k \leq p-1} \frac{1}{j^m k^m} \right) \\
&= H_{p-1,3m} + 3 \sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^{2m} k^m} + \frac{1}{j^m k^{2m}} \right) + 6 \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^m j^m k^m}.
\end{aligned}$$

As $H_{p-1,m} \equiv 0 \pmod{p}$, from the above we obtain

$$\begin{aligned}
\sum_{k=1}^{p-1} \frac{H_{k,m}^2}{k^m} &\equiv H_{p-1,3m} + \sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^{2m} k^m} + \frac{2}{j^m k^{2m}} \right) \\
&\quad - \frac{H_{p-1,3m}}{3} - \sum_{1 \leq j < k \leq p-1} \left(\frac{1}{j^{2m} k^m} + \frac{1}{j^m k^{2m}} \right) \\
&= \frac{2}{3} H_{p-1,3m} + \sum_{1 \leq j < k \leq p-1} \frac{1}{j^m k^{2m}} \pmod{p^2}.
\end{aligned}$$

Thus, by (3.1), (3.3) and the congruence $H_{p-1,3m} \equiv 0 \pmod{p}$, we immediately get (1.3).

Below we assume that $p > 3m + 1$. Adding (3.2) and (3.4) we obtain

$$\begin{aligned}
2 \sum_{1 \leq j < k \leq p-1} \frac{1}{j^m k^{2m}} &\equiv pm B_{p-1-3m} \left(-\frac{3}{3m+1} + \frac{\binom{3m}{m}}{(m+1)(2m+1)} \right) \\
&= \frac{pm}{3m+1} \left(\frac{\binom{3m+1}{m}}{m+1} - 3 \right) B_{p-1-3m} \pmod{p^2}.
\end{aligned}$$

Note also that

$$H_{p-1-3m} \equiv p \frac{3m}{3m+1} B_{p-1-3m} \pmod{p^2}.$$

Therefore

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{H_{k,m}^2}{k^m} &\equiv \frac{2}{3} \times p \frac{3m}{3m+1} B_{p-1-3m} + \left(\frac{\binom{3m+1}{m}}{m+1} - 3 \right) \frac{pm/2}{3m+1} B_{p-1-3m} \\ &= \left(\frac{\binom{3m+1}{m}}{m+1} + 1 \right) \frac{pm/2}{3m+1} B_{p-1-3m} \\ &= \left(\binom{3m+1}{m-1} + \frac{m}{2} \right) \frac{pB_{p-1-3m}}{3m+1} \pmod{p^2}. \end{aligned}$$

This proves (1.4).

So far we have completed the proof of Theorem 1.2. $\square$

REFERENCES

- [C] S. W. Coffman, *Problem 1240 and Solution: An infinite series with harmonic numbers*, Math. Mag. **60** (1987), 118–119.
- [G] H. W. Gould, *Combinatorial Identities*, Morgantown Printing and Binding Co., 1972.
- [IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory* (Graduate texts in math.; 84), 2nd ed., Springer, New York, 1990.
- [M] R. Meštrović, *A proof of a conjecture by Sun on congruence for harmonic numbers*, preprint, arXiv:1108.1171. <http://arxiv.org/abs/1108.1171>.
- [SW] J. Sondow and E. W. Weisstein, *Harmonic Number*, MathWorld—A Wolfram Web Resource, <http://mathworld.wolfram.com/HarmonicNumber.html>.
- [S] Z. H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (2000), 193–223.
- [Su] Z. W. Sun, *Arithmetic theory of harmonic numbers*, Proc. Amer. Math. Soc. **140** (2012), 415–428.
- [ST] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [W] J. Wolstenholme, *On certain properties of prime numbers*, Quart. J. Math. **5** (1862), 35–39.
- [ZS] L. L. Zhao and Z. W. Sun, *Some curious congruences modulo primes*, J. Number Theory **130** (2010), 930–935.